

# **Bubbles Navigating through Networks of Microchannels**

## **( Supplementary Information )**

Wonjae Choi, Michinao Hashimoto, Audrey K. Ellerbee, Xin Chen, Kyle J. M. Bishop, Piotr Garstecki<sup>†</sup>, Howard A. Stone<sup>‡</sup>, and George M. Whitesides<sup>\*</sup>

Department of Chemistry and Chemical Biology, Harvard University,

12 Oxford Street, Cambridge, MA 02138

<sup>†</sup> Institute of Physical Chemistry, Polish Academy of Sciences, Kasprzaka 44/52, 01-224

Warszawa, Poland

<sup>‡</sup> Department of Mechanical and Aerospace Engineering, Princeton University, Princeton, NJ

08544


\*Corresponding Author

Email Address: [gwhitesides@gmwhgroup.harvard.edu](mailto:gwhitesides@gmwhgroup.harvard.edu)

**Table S1.** Occupancy of channels by bubbles and the selection of the path at the first intersection of the network shown in Figs. 2, 4, 5.

	Occupancy of channel (# of bubbles)					The path an incoming bubble selects
	A	B	C	D	E	
(1)	0	0	1	1	1	Right (A)
(2)	0	0	1	0	1	Right (A)
(3)	0	0	2	1	0	Right (A)
(4)	1	0	1	0	2	Right (A)
(5)	0	0	1	1	0	Left (B CD E)
(6)	1	0	0	1	1	Left (B CD E)
(7)	1	0	1	0	1	Left (B CD E)
(8)	1	0	1	1	0	Left (B CD E)
(9)	1	0	1	0	0	Left (B CD E)
(10)	1	0	2	0	1	Left (B CD E)
(11)	2	0	1	0	0	Left (B CD E)

**Note:** The data set (3) and (5) determined the lower and upper bound of the resistance of a bubble, respectively.



**Figure S1.** The flow field of a single-phase Newtonian fluid in a point-symmetric two-path network. The Reynolds number is set to be zero so that the flow is Stokesian whose flow field can be calculated by solving Equations 1 and 2. The viscosity and density of the hypothetical fluid are set to  $1 \text{ (kg m s}^{-1}\text{)}$  and zero, and the pressures at the inlet (top) and outlet (bottom) are 1 (Pa) and zero. Obviously the flow field is point-symmetric as well, making the volumetric flows across the A-A' and B-B' cross-sectional areas to be identical.