

A macroscopic device described by a Boltzmann-like distribution†

Cite this: *Soft Matter*, 2013, 9, 4480

Simon Tricard,‡ Claudiu A. Stan,‡* Eugene I. Shakhnovich
and George M. Whitesides*

Equilibrium thermodynamic phenomena such as the Maxwell–Boltzmann distribution of molecular velocities are rare in systems of macroscopic particles interacting by mechanical collisions. This paper reports a system composed of millimeter-sized polymer objects that under mechanical agitation exhibits a “discretization” of the configurations of the system, and has a distribution of the probabilities of these configurations that is analogous to a Boltzmann distribution. The system is composed of spheres and a three-link chain on a bounded horizontal surface, shaken with an aperiodic but not completely random horizontal motion. Experiments were performed at different strengths of agitation (quantified by the frequency of agitation, f , at constant amplitude) and densities of spheres (quantified by the filling ratio, FR). The chain was typically found in one of three conformations – extended, single folded, and double folded – because, under collisions with the spheres, adjacent links were stable mechanically only when fully extended or fully folded. The probabilities of the different conformations of the chain could be described by a Boltzmann distribution in which the “temperature” depended on f and the “energies” of conformations on FR. The predictions of the Boltzmann formula using empirically determined “temperatures” and “energies” agreed with measurements within two experimental standard deviations in 47 out of 48 experiments.

Received 16th October 2012
Accepted 6th March 2013

DOI: 10.1039/c3sm27385g

www.rsc.org/softmatter

Introduction

A ubiquitous concept in statistical mechanics is the modeling of a gas by solid spherical particles colliding elastically in a box. This model is the first that most students encounter when they study statistical mechanics, and it explains clearly how thermodynamic properties (pressure, temperature, entropy, and many others) emerge as a result of molecular motions. The value of this model, aside from the validity of its analytical predictions, resides in its physically intuitive nature: it is built on basic concepts such as solid spherical particles and walls with precisely defined positions and velocities, perfectly elastic collisions, and so on.

It would be interesting to build a physical macroscopic model of a classical gas, even if only as a tool to teach statistical mechanics, but such a model is not (rigorously) possible

because macroscopic systems are dissipative. In order to maintain motion in a macroscopic system, one needs to supply energy to the system, and driven systems are not in thermodynamic equilibrium. External driving is a technique often used in the field of granular physics, which studies mechanical ensembles of moving macroscopic particles whose intrinsic thermal motion is too small to be observed.¹ In most cases, mechanically agitated systems of macroscopic particles do not exhibit thermodynamic equilibrium properties, such as a Maxwell–Boltzmann distribution of particle velocities.^{2,3}

It is nevertheless possible, though not trivial, to build physical systems of macroscopic particles whose motions mimic accurately the thermal agitation in a molecular gas. Examples of these systems are still rare,^{4,5} and pose interesting questions: can characteristics of equilibrium thermodynamic behavior be observed in a dissipative driven system? Do non-equilibrium systems that obey the laws of thermodynamics exist – and why?

Here we report a model macroscopic system (made from polymer beads shaken on a horizontal flat surface) that we designed to mimic a fundamental statistical-mechanical problem: a system with discrete energy levels in thermodynamic equilibrium with a thermal bath. More specifically, we investigated the behavior of a system analogous to a molecule with discrete vibrational levels immersed in a monoatomic buffer gas; the concentration of molecules is much lower than that of

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA. E-mail: claudiu.stan@alum.mit.edu; gwhitesides@gmwhgroup.harvard.edu

† Electronic supplementary information (ESI) available, containing: (1) additional information about the experimental setup; (2) all distinct folding shapes for each conformation of the chain; (3) values of the $E_{j,MA}/T_{MA}$ parameters for distinct experiments; and (4) a comparison between experimental measurements and the predictions of the Boltzmann model based on thermodynamic degeneracy values. See DOI: 10.1039/c3sm27385g

‡ These authors have contributed equally to this work.

the buffer gas and the molecule–molecule interactions are practically inexistent. The statistical–mechanical behavior of the molecule arises after many random collisions with atoms from the buffer gas.

Our model system is made of three polymer cylinders joined by flexible links (the “molecule”), immersed in an ensemble of free-rolling spheres (the “atoms”), and shaken continuously. The first surprising behavior of the chain was an approximate “discretization”: though the links are continuously bendable, the chain was found most of the time in only three spatial conformations (extended, and folded in two different ways) to which we were able to assign distinct “energies”. The second surprising behavior was thermodynamic-like statistical behavior: the probability of the chain being in one of the conformations was approximately described by a Boltzmann-like distribution.

Background

Physical-model simulations

Physical-model simulations can be used to study systems whose behavior is too complex to be modeled based on analytical calculations. Physical models are less common than numerical simulations, because numerical simulations, which benefit from the constant improvements in computational power and the sophistication of software, are often less expensive. There are nevertheless cases in which the phenomena under study are too complicated to rely exclusively on computational simulations. Physical models are still being built to study environmental phenomena such as soil erosion,^{6,7} atmospheric flows over rural terrains,⁸ physiological systems and processes,^{9,10} and technological procedures in metallurgy^{11,12} and drilling.¹³

We are interested in physical-model simulations for a different reason: we want to build models that can be perceived with our senses (sight, primarily) of phenomena that can only be observed with the aid of special tools, or for which direct component-level observations are not possible. We expect these physical models to guide our intuition about the phenomena that we simulate, and lead to discovery of new ones, because the human mind is specialized in interpreting information acquired by our senses.

As part of our previous explorations of self-assembly and of complexity, we have used ensembles of millimeter-sized objects to create several macroscopic models of molecular phenomena. The most ubiquitous phenomenon that we simulated was crystallization, which we could reproduce at the macroscopic scale using hard objects interacting by capillary¹⁴ and electrostatic^{15–18} forces. The capillary and electrostatic interaction potentials between objects were not identical to intermolecular potentials, but we could build a system with a particle interaction potential close in shape to intermolecular potentials, by using soft deformable objects that interacted by capillary forces; we used this system to simulate indentation fractures in crystals.¹⁹ We also simulated the dynamic behavior of polymers using mechanically-agitated macroscopic beads arranged on a string, and we observed bending and folding phenomena in systems in which the beads either interacted *via* electrostatic forces²⁰ or did not.²¹

Granular matter as a physical model of molecular thermodynamic behavior

Granular matter is composed of distinct particles; if the particles move independently from each other, a state that we will refer to as dynamic, the primary mechanism of interaction between particles is through collisions. Dynamic granular ensembles seem analogous to molecular gases, and attempts have been made to understand the properties of granular matter using the formalism of equilibrium thermodynamics.²² Parameters such as the granular temperature²³ have been used as surrogates for the thermodynamic temperature and proved useful in understanding the behavior of granular systems. The granular temperature T_G is equal to the average of the *fluctuations* of the kinetic energies of the particles. The velocity of the overall motion (the coarse-grained velocity of particles, \bar{v}_c) must be subtracted from the velocity of particles, v , as shown in eqn (1), where m_G is the mass of the particles.

$$T_G = \frac{m_G}{2} \overline{(v - \bar{v}_c)^2} \quad (1)$$

Granular matter in motion is a dissipative system, because collisions between particles are not elastic; external driving forces are required to maintain the movement of particles. The driving must be uniform, in the sense that all particles experience the same average driving force, to enable a thermodynamic distribution of the properties of particles; if the driving is not uniform, the granular temperature is highest near the driving source and decreases away from it.^{24,25} Though necessary, uniform driving is not sufficient to create a physical model of a thermodynamic system at equilibrium. In two-dimensional, vertically-vibrated submonolayer experiments, the distribution of velocities of granular particles is not the Gaussian curve characteristic of a Maxwell–Boltzmann distribution.^{2,3}

Experimental systems in which macroscopic objects behave similarly to molecules under thermal agitation have been reported previously.^{4,5} The key to the success of these systems seems to be the randomization of the driving forces. In a first example, two layers of spheres, gravitationally-segregated because of their different densities, were agitated vertically by a bottom plate;⁴ the two-dimensional velocities of particles from the top (but not the bottom) layer had a Gaussian distribution. In a second example, a granular mixture of spheres with two different sizes was sheared in a Couette cell (system thickness \sim five monolayers, driven from both bounding surfaces); the diffusivity and the mobility of tracer particles embedded in this system followed a Stokes–Einstein-type relation which was used to calculate the granular temperature.⁵

While eqn (1) represents probably the most intuitive way in which a granular temperature can be defined (*i.e.*, the temperature is a measure of the random kinetic energy of particles), calculating T_G from a Stokes–Einstein-type relation illustrates how a statistical-mechanical phenomenon (*i.e.*, the proportionality of diffusivity and mobility) can be used to define thermodynamic-analogous parameters for granular matter. Here we show that a statistical-mechanical relation, the Boltzmann distribution, can be used to define and calculate granular variables analogous not only to the temperature, but also to the

values of discrete energy levels and to statistical-mechanic degeneracies.

Experimental design

MecAgit: a horizontally-driven, two-dimensional granular system

Our experimental system (Fig. 1(a)) consists of a flat horizontal surface on which we shake millimeter-sized objects within a region bounded by vertical walls. Compared to other horizontally shaken granular systems, ours is distinguished by its pseudo-random agitation motion,²⁰ which is a combination of orbital shaking with randomly timed “kicks”; a detailed description of the pseudo-random agitation scheme can be found in the ESI.† To tune the intensity of agitation, we varied the frequency of the orbital shaker, f .

We prepared a circular mixing area with a diameter of 0.48 m using an aluminum rim, and we covered the area inside the rim with paper to generate an area with a constant friction coefficient on which the objects would roll, but not slide, when the plate was agitated. To avoid any possible electrical charging by contact electrification within the experimental setup, we maintained a relative humidity of more than 60% RH using a humidifier connected to the enclosed space above the plate.¹⁵

In all experiments, we filled the mixing area with simple (spheres) and composite (cylinders connected by a string) polymeric objects, and shook them with a pseudo-random motion to simulate molecular phenomena. We will refer to this method of simulating molecular phenomena as “mechanical agitation”, or MecAgit.²¹ The characteristics of MecAgit models are their two-dimensionality, the millimeter-size of the objects, the pseudo-random agitation, the possibility to design composite objects such as beads-on-a-string, and the control of long-range electrical interactions between objects by using different values of the relative humidity to suppress or allow contact electrification.

Designing a physical model of a canonical ensemble

The canonical ensemble is one of the most useful statistical-mechanical concepts for predicting the thermal behavior of a system. The whole system is composed of a smaller system (therefore referred to as simply “system”) that has a range of possible macrostates whose total energy E_i is known, and a thermal bath in which the system is immersed. The probability P_i of the system being in a macrostate with energy E_i depends exponentially on the absolute temperature T , and the probabilities are given by the Boltzmann distribution (eqn (2)), in which k_B is the Boltzmann constant, and the degeneracy g_i is the number of distinct microstates having the same energy E_i . The normalization factor is the partition function $Z(T)$, given by eqn (3), where the sum is conducted over all possible energies of the system.

$$P_i = \frac{1}{Z(T)} g_i e^{-\left(\frac{E_i}{k_B T}\right)} \quad (2)$$

$$Z(T) = \sum_i g_i e^{-\left(\frac{E_i}{k_B T}\right)} \quad (3)$$

Eqn (2) is general and can describe the behavior of a system of components which interact, because the total energy of the system includes the energy of interaction. The Boltzmann distribution also applies to other groups of microstates if the interactions between components are constant and the energies E_i can be written as a sum of terms corresponding to different degrees of freedom of the system. For example, in an ideal gas the Boltzmann distribution applies to the kinetic energy of the particles as well; this particular Boltzmann distribution was derived from the general one by averaging out the internal degrees of freedom of the particles. Here, we use the term “Boltzmann distribution” to refer to both the general case and to derived distributions. For derived distributions, E_i is the type of energy (e.g. kinetic) associated with the considered degrees of freedom, and g_i the number of microstates having the energy E_i .

To simulate a canonical ensemble using MecAgit, we needed a “thermal bath” and a “system”. The “thermal bath” was composed of free-rolling poly(methyl methacrylate) (PMMA) spheres with a diameter of 6.35 mm. We quantified the density of spheres in the system using the filling ratio (FR) parameter,




Fig. 1 (a) Experimental apparatus. Polymer spheres and a chain with flexible links are shaken with an aperiodic motion on a horizontal surface. (b) The three observed conformations for the chain: extended (C1), partially folded (C2), and fully folded (C3).

defined as the ratio between the number of spheres in the system and the number of spheres required to fill completely the mixing area in a single layer with hexagonal packing.

The “system” (Fig. 1(b)) was a chain composed of three Nylon cylinders (6.35 mm diameter, 14 mm length) connected by flexible links (14 mm length) between cylinders. The chain could assume three folding conformations (Fig. 1(b)): (i) C1: unfolded, with none of the cylinders in contact and extended, (ii) C2: partially folded, with two cylinders in contact, and (iii) C3: fully folded, with all three cylinders folded together. There are two conformations C3, with the middle cylinder either at the center or at the periphery of the folded chain. We used only one chain to suppress chain–chain interactions and to observe a Boltzmann-like distribution of the chain conformations. When multiple chains were shaken on the table, they interacted strongly and we could not observe the discretization of chain conformations.

The experiments consisted of measurements of the probability of conformations as a function of f (the orbital shaking frequency) and FR (the filling ratio). We recorded images of the system with a photo camera every 30 seconds for up to 30 minutes of continuous agitation, and we determined the type of conformation for each photograph using automated image analysis. We have focused on the “internal degrees of freedom” of the chain, ignoring translation. All measurements were performed during shaking and after the shaken system reached a steady state, because, similarly to the reasoning used in previous studies of dynamic granular systems, we regarded the shaking frequency as a measure of “temperature”; the steady-state system of spheres shaken at constant frequency thus had a constant “temperature”.

Measuring the probabilities of all conformations represents a statistical-mechanical approach and justifies the use of the canonical ensemble formalism²⁶ to describe our system, which is composed of only one particle (the chain) and thus is not thermodynamic. Another justification for treating the chain as a canonical ensemble with one particle is that the energy of interactions between the chain and the spheres is much smaller than the total energy of the chain; the energy of the chain, which depends on its conformation, represents a meaningful parameter because the fluctuations of its value are small during the timescale of interactions with the spheres. Since the spheres are less massive than the chain, a single collision between a sphere and the chain does not transfer enough energy to the chain to cause a transition between different conformations. We evaluated that the minimum number of collisions between individual spheres and the chain that caused a change in the conformation of the chain was on the order of 100, because (i) transitions between conformations took several seconds, (ii) the chain interacted on average with around ten spheres, and (iii) the characteristic frequency of interactions was on the order of the frequency of random “kicks” to the table, fixed at 4 Hz in all experiments. Between these transitions, the conformation of the chain was stable for periods on the order of tens of seconds.

In determining the statistical-mechanical properties of the system by temporal averaging of each conformation, we assumed implicitly that our canonical ensemble model is

ergodic, but we could not test this assumption using our setup. A test of ergodicity would require agitation of multiple chains at the same time, such that the temporal average of the conformation of one chain could be compared with the average conformation of all chains; also, the density of chains in the system must be low enough to prevent collisions between chains. The shaking table required for an ergodicity test would be several meters in diameter, and thus impractical to build.

Results

During agitation, the motion of the chain was caused by two driving forces: (i) friction with the agitating surface, and (ii) collisions with the PMMA spheres. Due to the design of our apparatus, the pseudo-random agitation engaged both the spheres and the chain, and was thus a uniform driving force. The mechanical driving of the chain was further randomized by collision with the spheres. The randomness of the overall driving of the chain and the dependence of conformation probabilities on the frequency of agitation suggested that the statistics of the conformations of the chain would be analogous to the statistics of an equilibrium thermal system.

To demonstrate that the statistics of our system was analogous to a Boltzmann distribution, we used this reasoning: (i) we assumed first that the statistics of the system was described by a Boltzmann distribution (eqn (2)), (ii) we calculated the parameters of the Boltzmann distribution, *i.e.* the “energies” of the three conformations of the chain, and the “temperature” of the chain, for each experiment, and (iii) we concluded that the statistical behavior of the system was analogous to that of a canonical ensemble, because we could fit with good accuracy the probabilities of configurations using a small number of “energy” and “temperature” parameters. Our reasoning assumed that (i) our system is ergodic; (ii) the chain–sphere interactions do not have to be included explicitly in the analysis because they were constant at a given set of f and FR values; we regarded them as a source of weak stochastic interactions that enabled the folding of the chain and that could be averaged out in the Boltzmann distribution; and (iii) the spheres shaken at constant frequency provide a “constant-temperature bath”. We could not demonstrate experimentally the validity of these separate assumptions, but we regard them as reasonable hypotheses that were collectively validated by a good fit of the Boltzmann model.

The “discretization” of the shape of the chain during agitation

We found that during agitation, the chain assumed the shape of one of the conformations C1, C2, or C3 in 93% of all photographs that we recorded. This “discretization” of shape to three distinct conformations during agitation was caused by the collisions with the spheres and by the particular relations between the sizes of the spheres, of the cylinders, and the mean free path of the spheres in our system. We can understand this phenomenon by considering a chain with two cylinders joined by a weakly elastic wire, colliding with an ensemble of spheres (with sizes comparable with those of cylinders) whose motion

does not have a directional bias. If the angle between the two cylinders is 180° (*i.e.* an extended conformation), the two sides of a single cylinder are impacted (on average) at the same rate by spheres, and though the angle between cylinders might fluctuate, there is no net “sphere pressure”²⁷ to cause the folding of the chain. If, however, the angle between cylinders is less than 180° and the mean free path of spheres is larger than the length of the cylinder, the sides of a cylinder that face the inside of the angle are partially shielded from sphere collisions by the other cylinder; in this case, there is a net “sphere pressure” that tends to reduce the angle and to fold the chain.

Depending on the stiffness of the link, there is a critical angle between the cylinders at which the tension of the link is balanced by the imbalance in sphere pressure. If the angle is larger than the critical angle, the elasticity of the link opens the chain; if the angle is smaller, the “sphere pressure” folds the chain. The two-cylinder chain is therefore a bistable system, with two stable equilibrium positions at cylinder angles of 0° and 180° , and one unstable equilibrium position at the critical angle. Our three-cylinder chain has three stable conformations; under the effect of fluctuations in the rate of collisions with spheres, the three-cylinder chain will be, for the majority of time, in one of its three conformations. The shape of the chain is thus “discretized”, in the approximate sense that only three conformations are *observed*.

The simple model we presented here to explain the predominance of C1, C2, and C3 conformations does not take into account biases in the motion of spheres, nor the fact that in our experiments, FR had values between 0.5 and 0.9, and the spheres therefore constituted a liquid-like rather than a gas-like system. We nevertheless believe that this model explains correctly the origin of “discretization” in our macroscopic system.

The dependence of the “energy levels” on the filling ratio

Because the links between cylinders are elastic, the folded conformations C2 and C3 have a higher potential energy than the open conformation C1. The mechanical elastic energy stored in once (C2) or twice (C3) folded links is not identical to the “energies” of conformations C2 and C3 that are the parameters of a Boltzmann-like distribution. An important characteristic of our system is that the statistic-mechanical “energies” associated with the three conformations of the chains depended on the filling ratio of the spheres. This dependence can be understood qualitatively using either of the two physical arguments below.

The mechanical energy needed to fold the chain when it is immersed in an ensemble of colliding spheres is less than the mechanical energy needed to fold the chain in the absence of collisions with spheres. When spheres are present, the chain needs to bend only as far as the critical angle for unstable equilibrium; past this point, the “sphere pressure” will complete the folding of the chain. At larger filling ratios, the surface density of spheres is larger, and thus the “sphere pressure” is larger; this fact brings the critical angle closer to 180° and reduces the energy needed for folding. As the filling ratio

increases, the differences in “energy” between the three conformations are thus reduced, and, as we will show later, the “energies” of the three conformations can become approximately equal.

An alternate explanation of the dependence of folding energies on FR is based on the analogy between the three-cylinder chain and a polymer molecule. Flexible polymer molecules whose monomer units do not interact assume a coiled shape with an average end-to-end length that is smaller than the contour length of the polymer.²⁵ Coiling is a statistical phenomenon caused by thermal agitation (*i.e.*, collisions with solvent molecules), and leads to measurable entropic forces that compress stretched polymers to the coiled conformation.^{28,29} In our system, sphere collisions tend to fold the three-cylinder chain, and the folding force is larger at higher filling ratios because the rate of collisions is larger. For the conditions of our experiments, these folding forces are “weaker” than the mechanical elasticity of the chain, and they reduce the *effective* stiffness of the chain, and thus the differences between the energies of the three conformations, by a degree that increases with the filling ratio.

The dependence of the conformation of the chain on FR and f

Fig. 2 shows the probabilities P_i of all conformations C_i ($i = 1, 2, 3$). We measured P_i for all combinations of FR values of 0.0, 0.5, 0.6, 0.7, 0.8, and 0.9, with f values of 100, 120, 140, and 160 rpm. Orbital agitation at frequencies below 80 rpm made spheres move collectively without any sphere–sphere collisions, corresponding to a granular temperature $T_G = 0$. At 80 rpm the spheres began to collide, but the evolution of the system towards a steady state was very slow. For orbital agitation frequencies above 160 rpm, the weight of the plate was not sufficient to maintain all four supporting cables extended, and the plate did not remain horizontal at all times. With the exception of the measurements at FR = 0.0 (no spheres), we chose the FR values to cover the widest range of filling ratios for which all three conformations were observed; for FR < 0.5, the chain either remained in the extended state, or the plate could not be leveled well enough to avoid movement of the chain towards the edge of the mixing area.

In the absence of spheres, the chain stayed extended (conformation C1) at all times, and $P_1 = 1$ and $P_2, P_3 = 0$. From FR = 0.5 to FR = 0.8, as f was increased, conformation C1 (the lowest energy) became less probable, and conformation C3 (the highest energy) became more probable. A special case was encountered for a filling ratio of 0.9: within experimental uncertainty, the probabilities did not depend on f and $P_1 < P_2 < P_3$.

If we regard the frequency of orbital agitation as a variable analogous to the thermodynamic temperature, the dependence of the coiling state of the chain on f is analogous to the behavior of certain polymer molecules. In the case of FR = 0, we have shown previously²¹ that chains similar to the one we use here, and agitated without spheres, followed the worm-like chain model (WLC) which describes the coiling of polymers that have a finite stiffness. When spheres are present, they make the whole system analogous with a polymer molecule in a “poor”




Fig. 2 Probabilities, P_i , of the chain being in one of the conformations C_i (see Fig. 1(b)) as a function of the filling ratio FR and of the frequency of orbital shaker f . For filling ratios between 0.5 and 0.8, as f increases, the probability of the lowest-energy state C1 decreases, and that of the highest-energy state C3 increases, suggesting that f is related to the granular temperature of the system. The investigated values of f were the same for all conformations; for clarity, data for C1 and C3 were shifted slightly along the x -axis. The error bars are the expected statistical errors, assuming a Poissonian distribution of the number of times, n_i , in which conformation was observed during an experiment (relative error: $(n_i + 1)^{-1/2}$; n_i varied between 0 and 35 among all measurements).

solvent with a lower critical solution temperature (LCST).³⁰ Usually, polymers in solvents tend to have a more extended configuration as the temperature increases, but solvents with a LCST show the opposite dependence – the polymer molecules become more compact as the temperature increases. In our case, the chain becomes more compact as the orbital agitation frequency f increases.

Calculation of the degeneracy of chain conformations

The statistical-mechanical definition of degeneracies g_i in our system is the number of “microstates” of the chain that have the same folding conformation C_i . A direct counting of the “microstates” was not possible, because we could not define the “microstates” of the chain. Instead, we attempted alternate calculation methods.

Thermodynamic method. We calculated the degeneracy values of the three conformations based on the number of folded shapes that cannot be transformed into each other by two-dimensional rotations and translations. A transition between two folded shapes of the same conformation can only happen if

the chain assumes a different conformation during the transition. Fig. S1 in the ESI† shows all possible folded shapes: one for C1, two for C2, and four for C3. Identifying the number of folded shapes with the degeneracy values, we obtained $g_1 : g_2 : g_3 = 1 : 2 : 4$.

Empirical method. The thermodynamic method is related to the concept that transitions between states with same energy but different symmetries can only occur *via* intermediate transitions to states with different energies. The number of states with different symmetries is thus related to the network of possible transitions between all states of the system. The degeneracies can be calculated from the network of these transitions, or “reaction paths”,³¹ and it was shown that for molecular systems the calculation of degeneracies based on counting the “reaction paths” is equivalent to the thermodynamic method based on symmetry.³² Our system has characteristics not encountered at the molecular scale (*e.g.* friction) that may lead to degeneracy numbers different from those calculated by the thermodynamic method. For example, a given way of folding the chain (*i.e.*, the “reaction path” in our system) could have a different rate than the reverse process of unfolding.

Given the subtleties of applying models valid for molecules to our macroscopic system, we calculated the degeneracy numbers empirically, using the property of the Boltzmann distribution (eqn (2)) that, if the energies of conformations are identical, the probabilities of conformations are independent of temperature and proportional to their degeneracies.

To identify the experimental conditions for which the three chain conformations had the same “energy”, we assumed that the “temperature” of the system is a function of the orbital agitation frequency f , because the kinetic energies of the objects increased as f increased. At FR = 0.9, the probabilities were approximately independent of f and thus independent of the “temperature”. Assuming that for FR = 0.9 the “energies” of configurations had the same value, we scaled the average probability of conformations to get the empirical degeneracy values $g_1 : g_2 : g_3 = 1 : 9.5 : 60.5$.

The “optimal” degeneracy values. The quality of the fit between the Boltzmann-like model and experimental results depended on the degeneracy values we used. Using the empirical degeneracy numbers led to a much better agreement with the Boltzmann-like model than the thermodynamic degeneracy numbers. We also attempted to determine an optimal set of degeneracy numbers through numerical fitting, but we could not find one, because the fitting error function did not have a minimum at physically reasonable degeneracy numbers. We will report here only the Boltzmann-like parameters obtained using empirical degeneracies.

Modeling the statistics of chain conformations with a Boltzmann distribution

The dependence of the probabilities of conformations on f for filling ratios from 0.5 to 0.8 (Fig. 2) is similar to the dependence of the populations of a system with three energy levels on temperature. This similarity suggested that the probability of

C1, C2 and C3 conformations might be described mathematically by a Boltzmann distribution (eqn (2)) in which the energies E_i and the temperature T are functions of the experimental parameters f and FR.

Derivation of a Boltzmann-like mathematical model that predicts the probabilities of the three conformations of the chain requires the definition of “temperature” and “energies” as functions of the experimental parameters f and FR. One way to determine these dependencies is to assume that the “temperature” and “energies” are polynomial functions of both f and FR, and determine the polynomial coefficients by multivariate mathematical fitting of measured probabilities to the model. Because multivariate fitting might not converge, or lead to macroscopic T and E_i that are complicated functions of f and FR, we have chosen instead to use a model as simple as possible.

In our model, (i) T depends only on f because the “temperature” should be related to the frequency of agitation, and (ii) E_i depend only on FR. The second assumption represents a simplification from the argument that we made previously to explain why the energy levels depend on the filling ratio. If the energy levels depend on the rates of sphere impacts on the chain, they will depend on FR and f because the rate of collisions depends on both parameters. We nevertheless chose to neglect the probable dependence of E_i on f , because it made the model simpler, and as we will show later, the agreement between model and experiments was good.

With the assumption that the “temperature” depends on f but not on FR, and the “energies” depend on FR but not on f , the Boltzmann-like equation for probabilities of conformations becomes:

$$P_i(f, \text{FR}) = \frac{1}{Z(f, \text{FR})} g_i e^{-\left(\frac{E_{i,\text{MA}}(\text{FR})}{T_{\text{MA}}(f)}\right)}, \quad i = 1, 2, 3 \quad (4)$$

In eqn (4) we defined the “MecAgit temperature” T_{MA} , and the “MecAgit energies” $E_{i,\text{MA}}$, which are measured in the same energy units (e.g. Joules). This definition is equivalent to choosing the “Boltzmann-like constant” $k_{\text{B,MA}} = 1$ in the Boltzmann-like equation; therefore, eqn (4) does not contain a “Boltzmann-like constant”.

The fitting of measured probabilities to eqn (4) is difficult because of its nonlinearity. Instead, in order to determine $E_{i,\text{MA}}(\text{FR})$ and $T_{\text{MA}}(f)$, we first linearized eqn (4) by eliminating the partition function $Z(f, \text{FR})$. Eqn (5) shows that the ratio of populations of two conformations does not depend on Z :

$$\frac{P_j}{P_1} = \frac{g_j}{g_1} e^{-\left(\frac{E_{j,\text{MA}} - E_{1,\text{MA}}}{T_{\text{MA}}}\right)}, \quad j = 2, 3 \quad (5)$$

Because only the differences between energy levels are relevant in our analysis, we adopted the convention that $E_{1,\text{MA}} = 0$. With this convention, we used eqn (6) to calculate the values of $E_{2,\text{MA}}/T_{\text{MA}}$ and $E_{3,\text{MA}}/T_{\text{MA}}$ from the conformation probabilities P_j (see Fig. S2 in the ESI†). Using these values, we calculated T_{MA} as a function of f by averaging the data from experiments performed at different FR values, using eqn (7).

$$\frac{E_{j,\text{MA}}}{T_{\text{MA}}} = -\ln\left(\frac{P_j g_1}{P_1 g_j}\right), \quad j = 2, 3 \quad (6)$$

$$T_{\text{MA}}(f) = \frac{1}{n_{j,\text{FR}}} \sum_{j,\text{FR}} T_{\text{MA}}(120) \times \frac{E_{j,\text{MA}}/T_{\text{MA}}(120, \text{FR})}{E_{j,\text{MA}}/T_{\text{MA}}(f, \text{FR})} \quad (7)$$

In eqn (7), $n_{j,\text{FR}}$ is the number of non-zero P_2 and P_3 probabilities measured at the same f but different FR values ($n_{j,\text{FR}} = 7$ or 8 depending on P_3 ; we did not use experiments for which $P_3 = 0$ because they made $E_{3,\text{MA}}/T_{\text{MA}}$ infinite). We defined the units of temperature such that $T_{\text{MA}} = 1$ at $f = 120$ rpm.³³ Fig. 3(a) shows the dependence of T_{MA} values as a function of agitation frequency, which could be fitted with good accuracy to a power-law dependence: $T_{\text{MA}} = 6.3 \times 10^{-6} \times f^{2.5}$.

We calculated the energies $E_{2,\text{MA}}$ and $E_{3,\text{MA}}$ of the C2 and C3 conformations according to eqn (8), where n_f is the number of non-zero probability measurements carried at a given frequency f ($n_f = 3$ or 4). The dependence of the energy levels on FR, shown in Fig. 3(b), was approximately linear, and the linear dependence extrapolated to zero at $\text{FR} = 0.95 \pm 0.03$, in approximate




Fig. 3 The MecAgit temperature and energies T_{MA} and $E_{j,\text{MA}}$. (a) The values of T_{MA} as a function of the frequency of agitation f , calculated after choosing $T_{\text{MA}} = 1$ at 120 rpm. The symbols represent the T_{MA} data, and the line shows the best power-law fit of the data, $T_{\text{MA}} \sim f^{2.5}$. (b) The configuration energies $E_{2,\text{MA}}$ and $E_{3,\text{MA}}$ as a function of the filling ratio FR. The symbols represent the measurements, and the lines are linear fits weighted by the standard error of measurements. The error bars in all graphs represent one standard deviation of the measurements (seven or eight measurements for T_{MA} ; three or four measurements for $E_{j,\text{MA}}$).

agreement with our earlier assumption that the energy levels were degenerate at FR = 0.9.

$$E_{j,MA}(\text{FR}) = \frac{1}{n_f} \sum_f \frac{E_{j,MA}}{T_{MA}}(f, \text{FR}) \times T_{MA}(f) \quad (8)$$

To verify the validity of our analogy between our system and a canonical ensemble, we calculated the probabilities of conformations using the Boltzmann-like distribution (eqn (4)) with the fitted functions shown in Fig. 3 (power-law in f for T_{MA} , and linear in FR for $E_{j,MA}$), and compared them with experimental measurements. Fig. 4 shows that the Boltzmann-like statistics provided a good description of the behavior of the system; 38 of the 48 calculated probabilities fell within one standard deviation of measured probabilities, and 47 of 48 fell within two standard deviations. Overall, the Boltzmann-like statistics were most accurate at FR ≥ 0.6 and $f \geq 120$ rpm.

We believe that the disagreement between the model and experiments at $f = 100$ rpm is due to the fact that the spheres approximated less accurately a thermal bath than at larger frequencies. Up to agitation frequencies of 80 rpm, the spheres moved together and their motions were thus well determined; as f increased above 80 rpm the motions of the spheres became gradually more randomized, and it is possible that at $f = 100$ rpm the motions of spheres were not completely randomized.

It is possible that at FR = 0.5 the system was not in a regime in which the conformation of the chain can be accurately predicted by a Boltzmann distribution. Below FR = 0.5 the chain remained extended at all times, and we have shown previously that the shape of the chain at FR = 0.0 can be described by the WLC model.²¹ The fact that two distinct models described the shape of the chain at low and high filling ratios suggests that the chain undergoes a phenomenon analogous to a phase

transition for a filling ratio between 0.0 and 0.5. Such transitions between different coiling regimes, for example the coil-to-globule transition,³⁴ have been encountered in polymer-solvent systems.

Discussion

Among the macroscopic parameters in the Boltzmann-like model, the MecAgit temperature has the clearest relation to the microscopic temperature, qualitatively and quantitatively. The power-law proportionality between T_{MA} and $f^{2.5}$ reflects the concept that the temperature is a measure of kinetic energy, because the velocity of the shaking table is proportional to f . The significance of the energy levels $E_{i,MA}$ and of the degeneracy numbers g_i are also qualitatively clear: the energy levels arise due the mechanical bistability of the conformation of adjacent links in the chain (*i.e.*, either folded or extended) under random collisions with the spheres, and the degeneracy numbers reflect the fact that there are more possible “reaction paths” to the folding of a given configuration than to its unfolding.

The MecAgit energy levels $E_{j,MA}$ in our system are different from those of atomic systems because they depend strongly on FR, while the electronic levels of atoms, molecules, and crystals depend only weakly on pressure at ambient conditions; relative changes in the electronic energy levels comparable to those observed in our MecAgit system are possible, but they require pressures thousands of times larger than atmospheric pressure.³⁵

The Boltzmann-like statistics provided a significantly better fit with the measurements when they included the empirical degeneracy values. For comparison, Fig. S3 in the ESI† shows the predictions of the Boltzmann model based on thermodynamic degeneracies, in the same format used in Fig. 4 for the case of empirical degeneracies. We could not find a way to calculate a set of degeneracy values that would agree with the empirical degeneracies, but we believe that such a calculation would be dependent on the details of the system. The relative size of the chain and the spheres or the surface density of spheres might suppress some of the folding or unfolding mechanisms.

Conclusion

We have developed a new granular system, composed by a chain surrounded by free spheres in a two-dimensional configuration, which exhibited a phenomenon analogous to a microscopic system with discrete energy levels: the distribution of the probabilities of the different configurations of the system was mathematically analogous to a Boltzmann distribution. Our system exhibited a behavior characteristic to non-dissipative systems, although it was dissipative due to mechanical friction and inelastic collisions. We believe that the unusual thermodynamic-analogous behavior in our driven granular system is due to the uniform driving of all objects by the shaking surface, and due to the randomization of the motion of the chain. Randomization of the motion of the chain in our system was achieved by a combination of the aperiodic movement of the shaking surface with collisions between the chain and independently-moving spheres.




Fig. 4 Comparison between experimental measurements and the predictions of the Boltzmann-like statistics with experimentally determined “energies” and “temperatures”, for filling ratios $0.5 \leq \text{FR} \leq 0.8$. The symbols connected by thin dashed lines are the experimental measurements (also shown in Fig. 2), and the thick solid lines the predictions of the Boltzmann-like statistics. For clarity, the data sets for the C1 and C3 conformations have been shifted horizontally, slightly, from the measurement values, which are shown on the x-axis.

The system presented here is a first attempt to create a granular system which is analogous to a thermodynamic system with discrete energy levels, and provides a new parameter for the “temperature” of a granular system – the MecAgit temperature T_{MA} . A comparison between T_{MA} and the granular temperature T_G characteristic to the spheres would be useful in determining the applicability and advantages of using T_{MA} and T_G as surrogates for the thermodynamic temperature. For this study we could not measure the sphere velocities which are required to calculate T_G , because we used a photo camera for imaging; the maximum rate at which we could record images of the system was smaller than the rate of collisions between spheres and than the frequency of agitation.

The system and the results reported here are part of a larger program in which we aim to physically model microscopic phenomena at the macroscopic scale. One of the goals of this program is to build systems that enhance our intuition of microscopic phenomena, but such systems could be useful outside the lab as well. The MecAgit system is simple and inexpensive enough for implementation in classroom teaching, where physical models have been shown to engage students and increase their level of understanding.^{36,37}

Acknowledgements

We thank Dr Phillip W. Snyder, Dr Antoine Venaille, and Dr Meital Reches for insightful discussions, and Dr Elizabeth Maxwell for editing advice. This work was primarily supported by the US Department of Energy, Division of Materials Sciences & Engineering, under Award no. DE-FG02-OOER45852. GMW acknowledges salary support from the US Department of Energy, Office of Basic Energy Sciences, under Award no. DE-SC0000989.

Notes and references

- I. S. Aranson and L. S. Tsimring, *Rev. Mod. Phys.*, 2006, **78**, 641–692.
- J. S. Olafsen and J. S. Urbach, *Phys. Rev. Lett.*, 1998, **81**, 4369–4372.
- J. S. Olafsen and J. S. Urbach, *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 1999, **60**, R2468–R2471.
- G. W. Baxter and J. S. Olafsen, *Nature*, 2003, **425**, 680.
- C. M. Song, P. Wang and H. A. Makse, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 2299–2304.
- E. Védie, J. L. Lagarde and M. Font, *Earth Surf. Processes Landforms*, 2011, **36**, 395–407.
- V. Bagarello and V. Ferro, *Catena*, 2012, **95**, 1–5.
- H. Kozmar, *Environ. Fluid Mech.*, 2012, **12**, 209–225.
- I. R. Titze, S. S. Schmidt and M. R. Titze, *J. Acoust. Soc. Am.*, 1995, **97**, 3080–3084.
- I. Jurjevic, M. Rados, J. Oreskovic, R. Prijic, A. Tvrdeic and M. Klarica, *Coll. Antropol.*, 2011, **35**, 51–56.
- B. Wright, P. Zulli, Z. Y. Zhou and A. B. Yu, *Powder Technol.*, 2011, **208**, 86–97.
- D. G. Ma, W. Q. Chen and X. M. Che, *Can. Metall. Q.*, 2012, **51**, 31–38.
- C. C. Hird, Q. Ni and I. Guymer, *Geotechnique*, 2011, **61**, 993–999.
- N. B. Bowden, M. Weck, I. S. Choi and G. M. Whitesides, *Acc. Chem. Res.*, 2001, **34**, 231–238.
- B. A. Grzybowski, A. Winkleman, J. A. Wiles, Y. Brumer and G. M. Whitesides, *Nat. Mater.*, 2003, **2**, 241–245.
- G. K. Kaufman, M. Reches, S. W. Thomas, J. Feng, B. F. Shaw and G. M. Whitesides, *Appl. Phys. Lett.*, 2009, **94**, 044102.
- G. K. Kaufman, S. W. Thomas, M. Reches, B. F. Shaw, J. Feng and G. M. Whitesides, *Soft Matter*, 2009, **5**, 1188–1191.
- R. Cademartiri, C. A. Stan, V. M. Tran, E. Wu, L. Friar, D. Vulis, L. W. Clark, S. Tricard and G. M. Whitesides, *Soft Matter*, 2012, **8**, 9771–9791.
- V. R. Thalladi, A. Schwartz, J. N. Phend, J. W. Hutchinson and G. M. Whitesides, *J. Am. Chem. Soc.*, 2002, **124**, 9912–9917.
- M. Reches, P. W. Snyder and G. M. Whitesides, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 17644–17649.
- S. Tricard, E. Feinstein, R. F. Shepherd, M. Reches, P. W. Snyder, D. C. Bandarage, M. Prentiss and G. M. Whitesides, *Phys. Chem. Chem. Phys.*, 2012, **14**, 9041–9046.
- I. S. Aranson and L. S. Tsimring, *Granular Patterns*, Oxford University Press, Oxford, 2009.
- P. K. Haff, *J. Fluid Mech.*, 1983, **134**, 401–430.
- A. Kudrolli, M. Wolpert and J. P. Gollub, *Phys. Rev. Lett.*, 1997, **78**, 1383–1386.
- E. L. Grossman, T. Zhou and E. BenNaim, *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 1997, **55**, 4200–4206.
- D. C. McQuarrie, *Statistical Mechanics*, University Science Books, Sausalito, CA, 2000.
- S. Aumaitre, C. A. Kruelle and I. Rehberg, *Phys. Rev. E: Stat., Nonlinear, Soft Matter Phys.*, 2001, **64**, 041305.
- M. D. Wang, H. Yin, R. Landick, J. Gelles and S. M. Block, *Biophys. J.*, 1997, **72**, 1335–1346.
- C. G. Baumann, S. B. Smith, V. A. Bloomfield and C. Bustamante, *Proc. Natl. Acad. Sci. U. S. A.*, 1997, **94**, 6185–6190.
- M. Rubinstein and R. H. Colby, *Polymer Physics*, Oxford University Press, Oxford, UK, 2003.
- D. M. Bishop and K. J. Laidler, *J. Chem. Phys.*, 1965, **42**, 1688–1691.
- G. Ercolani, C. Pigué, M. Borkovec and J. Hamacek, *J. Phys. Chem. B*, 2007, **111**, 12195–12203.
- We chose $f = 120$ rpm for reference because it was the lowest agitation frequency for which we observed all three conformations at filling ratios ranging from 0.5 to 0.8.
- A. Y. Grosberg and A. R. Khokhlov, *Statistical Physics of Macromolecules*, AIP Press, Woodbury, NY, 1994.
- J. M. Besson, W. Paul and A. R. Calawa, *Phys. Rev.*, 1968, **173**, 699–713.
- A. M. Ingham and J. K. Gilbert, *Int. J. Sci. Educ.*, 1991, **13**, 193–202.
- A. E. Rivet and K. A. Kastens, *J. Res. Sci. Teach.*, 2012, **49**, 713–743.