

Synthesis of 3-Deoxy-D-manno-2-octulosonate-8-phosphate (KDO-8-P) from D-Arabinose: Generation of D-Arabinose-5-Phosphate using Hexokinase¹

Mark D. Bednarski², Debbie C. Crans, Robert DiCosimo³, Ethan S. Simon⁴,
Phillip D. Stein⁵ and George M. Whitesides^{*}


Harvard University
Department of Chemistry
Cambridge, MA 02138

Marilyn J. Schneider

Wellesley College
Department of Chemistry
Wellesley, MA 02181

Summary. Three enzymes (used in soluble form enclosed in a dialysis membrane) efficiently produce 3-deoxy-D-manno-2-octulosonate-8-phosphate (KDO-8-P, **4**) from D-arabinose (Ara, **1**) and phosphoenolpyruvate (PEP **3**); production of D-Ara-5-P **2** from Ara using hexokinase is the key step in the synthesis.

This letter reports a multi-gram enzymatic synthesis of 3-deoxy-D-manno-2-octulosonate-8-phosphate⁶ (KDO-8-P, **4**) from D-arabinose (Ara, **1**) and phosphoenolpyruvate (PEP, **3**). The key step in this synthesis is the production of the expensive intermediate D-arabinose-5-phosphate (Ara-5-P, **2**, ~ \$10⁴/mole) from D-arabinose (Ara, **1**) using hexokinase.⁷ KDO-8-P is a key intermediate in the biosynthesis of the lipopolysaccharide (LPS) region of gram-negative bacteria.^{8,9} Inhibitors of LPS biosynthesis are candidates in the design of antimicrobial pharmaceuticals.^{10,11} The availability of gram quantities of KDO-8-P may facilitate the search for new inhibitors. This synthesis may also be applicable to the synthesis of analogs of KDO-8-P.



Production of Ara-5-P **2** relies on two enzymes: hexokinase (E.C. 2.7.1.1) and pyruvate kinase (E.C. 2.7.1.40). Hexokinase catalyzes the generation of **2** and ADP from Ara **1** and a catalytic amount of ATP. In a coupled system, pyruvate kinase regenerates ATP from ADP, while converting the readily available PEP to pyruvate.¹² Aldol condensation of **2** with additional PEP using KDO-8-phosphate synthetase (E.C. 4.1.3.3), isolated from *E. coli* B, produces **1**.¹³ The enclosure of all three enzymes in a single dialysis membrane facilitates separation of the reaction products from protein and allows reuse of the catalysts.¹⁴ In our hands the enzymatic production of Ara-5-P is superior to the method based upon the reaction between D-glucosamine-6-phosphate and ninhydrin.¹⁵

Isolation of **4** is straightforward. Addition of barium chloride to the crude reaction mixture precipitates a mixture of barium salts containing primarily KDO-8-P and inorganic phosphate (P_i). Stirring the mixture with ion exchange resin (Dowex 50W, H⁺ form), followed by neutralization with LiOH and filtration, removes P_i as its insoluble lithium salt. Lyophilization of the filtrate followed by further purification by gel filtration (Biogel P-2, eluant: water) provides 90% pure **4** in 63% yield; PEP and KDO (trace) are also present. Ion exchange chromatography (AG 1-X2, eluant: 0 to 1.3 M $HCO_3^-NH_4^+$) removed residual PEP to give **4** in >95% purity (42% yield). Experimental procedures are provided below.

KDO-8-phosphate (4) from Ara (1). To a solution of $PEP \cdot K^+$ **12 3** (15.25 g, 74 mmol; pH 7) in water (250 mL) were added Ara **1** (9.00 g, 60 mmol) and ATP (3.68 g, 6 mmol) and the solution was adjusted to pH 7.6 by addition of 1 N NaOH. After addition of $MgCl_2 \cdot 6H_2O$ (1.76 g, 8.7 mmol) and 2-mercaptoethanol (0.1 mL), the reaction solution was placed in a 500-mL graduated cylinder containing a magnetic stirring bar, and was degassed with N_2 for 30 min. KDO-8-phosphate synthetase¹³ (50 U, 1 U = $1 \mu mol \cdot min^{-1} \cdot mg^{-1}$), pyruvate kinase (160 U) and hexokinase (16,000 U) were dissolved in 13 mL of the reaction solution, placed in a dialysis membrane (Spectrapor 2, MW cutoff 12,000-14,000, 25 mm) and added to the reaction vessel; N_2 was continuously bubbled through the solution. After 4 days, the enzyme-containing bag was removed and dialyzed twice against 50 mL of water. The combined reaction mixture and dialysates (350 mL) were transferred to a 2-L Erlenmeyer flask and $BaCl_2 \cdot 2H_2O$ (20 g, 82 mmol) was added, followed by 700 mL of acetone. The solution was stirred for 2 h and left overnight at 4 °C. Filtration of the resulting barium precipitate (which consisted primarily of P_i and KDO-8-P) yielded 26.4 g of a brown solid. Water (500 mL) was added to this solid followed by ion exchange resin (Dowex 50W, H⁺, 17 g) and the solution was stirred for 72 h. Neutralization of the filtrate with LiOH precipitated P_i . The filtrate was concentrated in vacuo to approximately 100 mL, and this solution was passed through a column of Biogel P-2 (100 x 4.5 cm, eluant: water). Fractions containing **4** were pooled¹⁶ and concentrated in vacuo to give 8.2 g of **4** (90% purity by 1H NMR spectroscopy; 63% yield); PEP (10%) and KDO (trace) were also present. Ion exchange chromatography (AG 1-X2, eluant: 0 to 1.3 M $HCO_3^-NH_4^+$) removed residual PEP. The fractions containing KDO-8-P were pooled and brought to pH 6 by

addition of Dowex 50W (H⁺ form) and the solution was filtered and concentrated in vacuo to give 5.41 g of **4** (>95% purity, 42% yield). The product was characterized by ¹H NMR, ¹³C NMR and by assaying for release of phosphate according to the procedure of Ray;⁶ treatment of **4** with acid phosphatase (E.C. 3.1.3.2) yielded KDO .

Arabinose-5-phosphate (2). To a solution of PEP⁻K⁺ **3** (11.0 g, 91%, 48 mmol; pH 7) in water (75 mL) in a 250-mL Erlenmeyer flask containing a magnetic stirring bar were added: Ara **1** (10.9 g, 72.8 mmol), ATP (1.29 g, 2.16 mmol) and MgCl₂·6H₂O (2.2 g, 10.8 mmol). The pH was adjusted to pH 7.6 with 1 N NaOH, the solution was degassed with N₂ for 30 min and 0.1 mL of 2-mercaptoethanol was added. Hexokinase (10,000 U) and pyruvate kinase (500 U) were dissolved in 4 mL of the reaction mixture and the enzyme solution was placed in a section of dialysis tubing . The enzyme-containing bag was added to the reaction flask which was then stoppered and agitated at 150 rpm on an orbital shaker at room temperature. Analysis by ³¹P NMR spectroscopy monitored the formation of **2**. After 4 days, 10,000 U of additional hexokinase in a dialysis membrane were added to the reaction mixture. After an additional 3 days, 10,000 U of hexokinase, 500 U of pyruvate kinase and 850 mg (4.2 mmol) of MgCl₂·6H₂O were added. A white precipitate, presumably the magnesium salt of P_i, formed during the reaction. After an additional 5 days, the enzyme-containing bags were removed and dialyzed twice against 50 mL quantities of water. To the combined reaction and dialysates was added 4 g of BaCl₂·2H₂O (in 1 g portions dissolved in approximately 10 mL of water) to precipitate remaining ATP and P_i. After each addition, the mixture was filtered through Celite and each filter cake was washed with approximately 30 mL of water. To the combined filtrates (300 mL) was added BaCl₂·2H₂O (13 g, 53 mmol) and then 900 mL of EtOH and the mixture was left at 4 °C overnight. The precipitate was collected by filtration, washed with 100 mL of EtOH:water (3:1, v:v) and dried in vacuo over P₂O₅ to give 18.5 g of the barium salt of **2** (85% purity, 89% yield). The spectral data matched that of a sample obtained from Sigma; PEP (5%) and dipyruvate (10%) were also present. Anal: Calcd. for C₅H₉O₈PBa: C, 16.43; H, 2.48; Ba, 37.58; Found: C, 16.33; H, 2.45; Ba, 37.17.

References

1. Supported by NIH Grant GM 30367.
2. American Cancer Society Postdoctoral Fellow, Grant No. PF-2762, 1986-88.
3. Standard Oil Co.(Ohio), Cleveland, Ohio.
4. DuPont Fellow 1986-87.
5. NIH Postdoctoral Fellow, Grant GM 10324, 1984-86.
6. This compound has been prepared enzymatically in milligram quantities from Ara-5-P and PEP: Ray, P. H. J. Bacteriol. **1980**, 141, 635.

7. Ara-5-P has not been prepared from Ara using hexokinase, although it is reported that Ara is a substrate for the enzyme ($v_{rel} \sim 0.02$): Sols, A.; De La Fuente, G.; Villar-Palasi, C.; Asensio, C. Biochem. Biophys. Acta **1958**, 30, 92.
8. Unger, F. M. Adv. Carbohydr. Chem. Biochem. **1981**, 38, 323.
9. Bacterial Endotoxin: Chemical, Biological and Clinical Aspects: Homma, J. Y.; Kanegasak, S.; Lüderitz, O.; Shiba, T.; Westphal, O., Eds.; Verlag Chemie: Weinheim, 1984.
10. Hammond, S. M.; Claesson, A.; Jansson, A. M.; Larsson, L.-G.; Pring, B. G.; Town, C. M.; Ekström, B. Nature **1987**, 327, 730.
11. Norbeck, D. W.; Kramer, J. B.; Lartey, P. A. J. Org. Chem. **1987**, 52, 2174.
12. Hirschbein, B. L.; Mazenod, F. P.; Whitesides, G. M. J. Org. Chem. **1982**, 104, 4458.
13. KDO-8-phosphate synthetase was isolated from E. coli B. cells (Grain Processing Company, 1600 Oregon St., P.O. Box 349, Muscatine, Iowa 52761) following the procedure of Ray⁶ with two modifications: a French Press was used for cell disruption and the enzyme fractions pooled after the DEAE Sephadex chromatography step (step 4) were stabilized with bovine serum albumin (50 mg/mL) and concentrated by lyophilization. Further purification was not necessary because the preceding steps removed most of the KDO-8-P phosphatase activity.
14. Bednarski, M. D.; Chenault, H. K.; Simon, E. S.; Whitesides, G. M. J. Am. Chem. Soc. **1987**, 109, 1283.
15. Volk, W. A. Methods Enzymol. **1966**, 9, 38.
16. Fractions containing **4** were identified by using the thiobarbituric acid assay (Karkhanis, Y. D.; Zeltner, J. Y.; Jackson, J. J.; Carlo, D. Anal. Biochem. **1978**, 85, 595) or by tlc analysis (silica gel, eluant: n-BuOH: AcOH:H₂O, 5:3:1).

(Received in USA 5 November 1987)