

Figure 1. Plots for solvent extraction of aqueous lithium ( $\circ$ ), sodium ( $\Delta$ ), potassium ( $\blacktriangle$ ), rubidium ( $\bullet$ ), caesium ( $\bullet$ ), silver ( $\square$ ), and thallium picrate ( $\blacksquare$ ) with 12-crown-4 in dichloromethane: (a) 1:1 or (b) 2:2 stoichiometry assumed;  $[12\text{-crown-4}]_{\text{org}}$  and  $[A^-]_{\text{aq}}$  refer to concentrations of 12-crown-4 in organic phase ( $\text{CH}_2\text{Cl}_2$ ) and of picrate ion in aqueous phase at the equilibrium, and  $D_M$  represents the distribution ratio of metal ion between organic/aqueous phases; for the derivation of equilibrium equations, see ref. 1.

enhanced by the bridging co-ordination of two picrate anions, since the lipophilicity of the complex, or the counter anion, to be extracted is one of the primary factors dominating extractability and cation selectivity in solvent extraction.<sup>3,7,8</sup> This does not mean however that there is no 1:1 or 1:2 complex formed in the aqueous phase, but rather indicates that only the highly lipophilic 2:2 complex, even though it may be a minor species in the aqueous phase, is extracted preferably into dichloromethane. Another point of interest is the cation selectivity. As a consequence of the different complex stoichiometry applied, selectivity inversion is actually observed for several cation pairs, e.g.  $\text{Ag}^+/\text{Tl}^+$ ,  $\text{Li}^+/\text{K}^+$ ,  $\text{Rb}^+/\text{Cs}^+$ , as indicated by the crossing plots in Figure 1.

The present result indicates that the ion-pair complex extracted into the organic phase from aqueous solution, though it shares some resemblance with the crystalline

complex,<sup>1</sup> possesses its own characteristic stoichiometry which is distinctly different from those observed in the homogeneous or solid phase. Thus the complex stoichiometry in solvent extraction is not a simple function of the size-fit concept but is influenced substantially by the lipophilicity of the ion-pair complex to be extracted. From the practical point of view, the different stoichiometry for the specific cation pair may be used as a tool for controlling cation selectivity.

Partial support of this work by the Ministry of Education, Science and Culture of Japan through a Grant-in-Aid for Scientific Research is gratefully acknowledged. Y. L. acknowledges the generous financial support of Ako Chemical Co. and Himeji Overseas Chinese Association for his stay at the Himeji Institute of Technology.

Received, 13th June 1989; Com. 9/02486G

## References

- 1 Y. Inoue, Y. Liu, F. Amano, M. Ouchi, A. Tai, and T. Hakushi, *J. Chem. Soc., Dalton Trans.*, 1988, 2735.
- 2 R. Hilgenfeld and W. Saenger, in 'Host Guest Complex Chemistry,' vol. 2, ed. F. Vögtle, Springer, Berlin, 1982, pp. 1 *et seq.*; M. Mercer and M. R. Truter, *J. Chem. Soc., Dalton Trans.*, 1973, 2469; D. L. Hughes, *ibid.*, 1975, 2374; J. D. Owen and M. R. Truter, *ibid.*, 1979, 1831.
- 3 F. de Jong and D. N. Reinhoudt, 'Stability and Reactivity of Crown Ether Complexes,' Academic, London, 1981, pp. 5 *et seq.*; H. K. Frensdorff, *J. Am. Chem. Soc.*, 1971, **93**, 600; R. M. Izatt, R. E. Terry, D. P. Nelson, Y. Chan, D. J. Eatough, J. S. Bradshaw, L. D. Hansen, and J. J. Christensen, *ibid.*, 1976, **98**, 7626; A. Hofmanova, J. Koryta, M. Brezina, and M. L. Mittal, *Inorg. Chim. Acta*, 1978, **28**, 73; Y. Takeda, *Bull. Chem. Soc. Jpn.*, 1979, **52**, 2501.
- 4 M. Dobler and R. P. Phizackerley, *Acta Crystallogr., Sect. B*, 1974, **30**, 2746, 2748. Interestingly, the ionophore antibiotic lasalocid A forms a 2:2:2 cation:ligand:water complex with sodium ions in the solid state, although only one water molecule bridges two sodium ions in this case; G. D. Smith, W. L. Duax, and S. Fortier, *J. Am. Chem. Soc.*, 1978, **100**, 6725.
- 5 R. Eiest and R. Weiss, *J. Chem. Soc., Chem. Commun.*, 1973, 678.
- 6 F. P. Van Remoortere and F. P. Boer, *Inorg. Chem.*, 1974, **13**, 2071; F. P. Boer, M. A. Neuman, F. P. Van Remoortere, and E. C. Steiner, *ibid.*, 1974, **13**, 2826.
- 7 Y. Takeda, in 'Host Guest Complex Chemistry,' vol. 3, ed. F. Vögtle, Springer, Berlin, 1984, pp. 1 *et seq.*
- 8 Y. Inoue, M. Ouchi, and T. Hakushi, *Bull. Chem. Soc. Jpn.*, 1985, **58**, 525.

## A New, General Method for the Determination of Binding Constants: Photoacoustic Titration of Dansylamide with Carbonic Anhydrase

Ahamindra Jain, Elaine M. Marzluff, John R. Jacobsen, George M. Whitesides, and Joseph J. Grabowski\*  
Department of Chemistry, Harvard University, 12 Oxford Street, Cambridge, MA 02138, U.S.A.

We have developed a new method utilizing the partitioning of energy between radiative and non-radiative pathways following photostimulation for the determination of binding constants in biochemical systems; application of this photoacoustic calorimetry technique to the study of the inhibition of carbonic anhydrase by 1-(dimethylamino)-5-naphthalenesulphonamide yields  $K_{11} = 1.7 (\pm 0.4) \times 10^6 \text{ mol}^{-1} \text{ dm}^3$ .

Photoacoustic calorimetry (PAC) has been applied by Peters,<sup>1</sup> Braslavsky,<sup>2</sup> Griller,<sup>3</sup> and others<sup>4</sup> to the determination of thermodynamic parameters and/or dynamics in a number of chemical systems, including the photodissociation of carbon monoxide from myoglobin,<sup>1b</sup> the displacement of heptane

from  $\text{Cr}(\text{CO})_5$ heptane by pyridine,<sup>1c</sup> and the measurement of bond dissociation energies.<sup>3</sup> We wish to report the first application of PAC to the determination of the affinity of a receptor for a ligand.

Carbonic anhydrase (bovine erythrocyte, mixture of isozy-




Figure 1. A typical photoacoustic titration curve for dansylamide-carbonic anhydrase. Circles represent experimental data; the curve is the calculated best fit, corresponding to  $K_{11} = 1.9 \times 10^6 \text{ mol}^{-1} \text{ dm}^3$ . (DNSA = dansylamide; CA = carbonic anhydrase;  $[\text{DNSA}]_0 = 2.04 \times 10^{-5} \text{ M}$ ;  $\lambda_{\text{exc}} = 337.1 \text{ nm}$ ; pulse energy  $\sim 20 \mu\text{J}$ ). Inset: A typical photoacoustic wave for dansylamide in water; the experimental baseline (signal for aqueous buffer alone) and the integration limits (used to obtain the signal amplitude) are also indicated.

mes) interacts with 1-(dimethylamino)-5-naphthalenesulphonamide (dansylamide) to afford a 1:1 complex.<sup>5</sup> The change in fluorescence of dansylamide observed upon its binding to carbonic anhydrase is the basis of an existing assay for the determination of binding constants of non-fluorescent inhibitors to this protein.<sup>6</sup> Herein we describe a new, general method which utilizes the difference in quantum yields for non-radiative relaxation exhibited by free and bound dansylamide. This difference in thermal energy deposition to the solution is used to monitor the extent of complex formation as a known concentration of inhibitor is titrated with protein.<sup>7</sup> Dansylamide, bound to carbonic anhydrase or free in solution, absorbs photons at 337.1 nm ( $\text{N}_2$  laser); the free ligand rapidly returns most of the absorbed energy to the solution as heat ( $\Phi_f$  is small;<sup>7</sup>  $f_h = 0.978 \pm 0.006$ ),<sup>8</sup> whereas the enzyme-bound molecule returns a smaller fraction of the excitation energy as heat ( $f_h \sim 0.3$ )<sup>8</sup> and emits the rest as fluorescence ( $\Phi_f = 0.84$ ).<sup>5</sup> Any non-radiative relaxation will produce a local temperature increase, which gives rise to an adiabatic, isobaric expansion, which in turn creates a pressure wave (acoustic wave) in solution. This pressure wave passes through the solution and is readily transmitted through the quartz wall of a standard cuvette and into a piezoelectric based transducer that is clamped to the outside of the cell. An

example of a detected photoacoustic wave for dansylamide in aqueous buffer is shown in the inset of Figure 1.

A PAC experiment to determine a binding constant is executed in the following manner: Degassed dansylamide in 0.3% methanol-20 mM aqueous  $\text{KH}_2\text{PO}_4$  buffer (pH 7.4) is placed in the sample cuvette, at room temperature, and photoacoustic waves generated from this sample upon irradiation by a short, low energy laser pulse ( $\leq 20 \mu\text{J}$ ) are digitized, collected, and averaged for 35 laser shots.<sup>§</sup> The equilibrium concentrations of inhibitor, protein, and their complex are then perturbed by increasing the protein concentration and a new set of data collected. Over the course of this titration of a fixed concentration of dansylamide, we observe a decrease in the amplitude of the photoacoustic signal that scales with the amount of carbonic anhydrase added. The data are analysed by integration of the first negative excursion of the photoacoustic waves (shaded area of Figure 1 inset) for each ratio of [carbonic anhydrase]-to-[dansylamide] and are then normalized to laser energy, which is also measured for each pulse during the experiment.<sup>9</sup>

The set of integrated, normalized photoacoustic signals from the titration are fitted by a non-linear least-squares routine to an expression for a 1:1 binding interaction. Typical data from one experiment are shown in Figure 1. The binding constant, obtained as the average from five separate experiments ( $\pm$  one standard deviation) is  $1.7 (\pm 0.4) \times 10^6$

<sup>†</sup> We also can and have obtained satisfactory data from the titration of carbonic anhydrase with dansylamide, an assay requiring less protein.

<sup>‡</sup> The variable  $f_h$  is the fraction of absorbed energy released as heat.

<sup>§</sup> No waveform shifts were observed upon titration, indicating that heat deposition is always occurring in the same time frame.

$\text{mol}^{-1} \text{dm}^3$ , in excellent agreement with a binding constant of  $2.1 (\pm 0.5) \times 10^6 \text{ mol}^{-1} \text{dm}^3$  obtained from non-linear fitting of data obtained by the fluorescence binding assay.<sup>10</sup> These values are in accord with previous measurements of this binding constant [ $4.2 (\pm 0.2) \times 10^6 \text{ mol}^{-1} \text{dm}^3$ ;  $5.0 \times 10^5 \text{ mol}^{-1} \text{dm}^3$ ].<sup>6</sup>

The PAC binding assay is both experimentally straightforward and versatile. Any system comprised of bound and free populations for which the non-radiative transitions following photostimulation are different for these species can in principle be examined by PAC. Changes in quantum yields for fluorescent and non-radiative processes or in time constants for energy deposition are two important parameters that the PAC technique can exploit, as well as changes in extinction coefficients for the species being studied. The method has obvious application where background fluorescence or light scattering interferes with measurements of fluorescence of the group of interest, or where luminescence is not easily monitored. We are now pursuing other systems for which PAC offers a unique solution to problems in molecular recognition.

We thank Carolyn R. Bertozzi for designing and fabricating our photoacoustic calorimeter. This work has been supported by the Harvard Materials Research Laboratory under National Science Foundation grant #DMR-86-14003. A.J. acknowledges a fellowship from Johnson & Johnson.

Received, 9th May 1989; Com. 9/01942A

References

- 1 (a) L. J. Rothberg, J. D. Simon, M. Bernstein, and K. S. Peters, *J. Am. Chem. Soc.*, 1983, **105**, 3464; (b) J. A. Westrick, J. L. Goodman, and K. S. Peters, *Biochemistry*, 1987, **26**, 8313; (c) G. K. Yang, V. Vaida, and K. S. Peters, *Polyhedron*, 1988, **7**, 1619; (d) K. S. Peters and G. J. Synder, *Science*, 1988, **241**, 1053.
- 2 S. E. Braslavsky, R. M. Ellul, R. G. Weiss, H. Al-Ekabi, and K. Schaffner, *Tetrahedron*, 1983, **39**, 1909.
- 3 T. J. Burkey, M. Majewski, and D. Griller, *J. Am. Chem. Soc.*, 1986, **108**, 2218.
- 4 (a) J. A. LaVilla and J. L. Goodman, *Chem. Phys. Lett.*, 1987, **141**, 149; (b) V. E. Anderson, H. Z. Cheng, G. J. Diebold, A. Mahmood, and D. A. Sweigart, *J. Am. Chem. Soc.*, 1987, **109**, 6191; (c) M. S. Herman and J. L. Goodman, *ibid.*, 1989, **111**, 1849; (d) T. Ni, R. A. Caldwell, and L. A. Melton, *J. Am. Chem. Soc.*, 1989, **111**, 457; (e) J. Rudzki-Small, J. J. Hutchings, and E. W. Small, in 'Fluorescence Detection III,' ed. E. R. Menzel, SPIE Proceedings, vol. 1054, Society of Photo-Optical Instrumentation Engineers, Bellingham, W.A., 1989, in the press.
- 5 R. F. Chen and J. C. Kernohan, *J. Biol. Chem.*, 1967, **242**, 5813.
- 6 G. S. Ponticello, M. B. Freedman, C. N. Habecker, P. A. Lyle, H. Schwam, S. L. Varga, M. F. Christy, W. C. Randall, and J. J. Baldwin, *J. Med. Chem.*, 1987, **30**, 591.
- 7 Y.-H. Li, L.-M. Chan, L. Tyer, R. T. Moody, C. M. Himei, and D. M. Hercules, *J. Am. Chem. Soc.*, 1975, **97**, 3118.
- 8 E. M. Marzluff, C. R. Bertozzi, J. R. Jacobsen, and J. J. Grabowski, manuscript in preparation.
- 9 J. J. Grabowski, J. D. Simon, and K. S. Peters, *J. Am. Chem. Soc.*, 1984, **106**, 4615.
- 10 A. Jain and G. M. Whitesides, unpublished results.

Reaction of Trifluoromethanesulphonyl Chloride with Alkenes catalysed by a Ruthenium(II) Complex

Nobumasa Kamigata,\* Takamasa Fukushima, and Masato Yoshida



Department of Chemistry, Faculty of Science, Tokyo Metropolitan University, Fukazawa, Setagaya-ku, Tokyo 158, Japan

The reaction of trifluoromethanesulphonyl chloride with alkenes in the presence of dichlorotris(triphenylphosphine)ruthenium(II) gives 1:1 adducts with extrusion of sulphur dioxide.

Recently, fluoroalkylated compounds have received much attention in medicine and material chemistry.<sup>1</sup> However, it is difficult to introduce a fluoroalkyl group into alkenes or aromatic compounds by ordinary alkylating methods *via* carbocationic intermediate because of the high electron-withdrawing character of fluorine. New methods of fluoroalkylation have been developed: *e.g.* fluoroalkylation of alkenes or aromatic compounds by perfluoroalkanoyl peroxide,<sup>2</sup> metal complexes of fluoroalkyl halides,<sup>3</sup> perfluorocarboxylic acids mediated by xenon difluoride,<sup>4</sup> fluoroalkyl halide with metal complex catalysts,<sup>5</sup> and electrochemical method with trifluoroacetic acid<sup>6</sup> have been used. However, these methods do not give satisfactory yields of the products, and often suffer from side reactions. We have already reported the reaction of trichloromethanesulphonyl chloride with alkenes in the presence of ruthenium(II) complex to give 1:1 adducts in good yields with evolution of sulphur dioxide.<sup>7</sup> During our systematic investigations on the reactions of sulphonyl chlorides with

alkenes in the presence of a catalytic amount of a ruthenium(II) complex, we found that the reaction of trifluoromethanesulphonyl chloride (1) with alkenes in the presence of a ruthenium(II) complex proceeded smoothly under mild conditions to afford the corresponding 1:1 adducts in high yields with extrusion of sulphur dioxide.

Typically, a solution containing (1) (2.0 mmol), styrene (2a) (5.0 mmol), and dichlorotris(triphenylphosphine)ruthenium(II) (0.02 mmol) in 4.0 ml of benzene was degassed and heated at 120 °C in a sealed tube for 16 h. Flash column chromatography on Florisil using benzene and then the gel permeation chromatography using chloroform gave 3-chloro-1,1,1-trifluoro-3-phenylpropane (3a) in 84% yield. Other metal catalysts,



Scheme 1